



Perforin Monoclonal Antibody (dG9 (delta G9)), eFluor™ 450, eBioscience™

Product Details	
Size	100 Tests
Species Reactivity	Human, Pig
Published Species	Human
Host/Isotype	Mouse / IgG2b, kappa
Recommended Isotype Control	Mouse IgG2b kappa Isotype Control (eBMG2b), eFluor™ 450, eBioscience™
Class	Monoclonal
Туре	Antibody
Clone	dG9 (delta G9)
Conjugate	eFluor™ 450
Excitation/Emission Max	405/445 nm
Form	Liquid
Concentration	5 μL/Test
Purification	Affinity chromatography
Storage buffer	PBS, pH 7.2, with 0.2% BSA
Contains	0.09% sodium azide
Storage conditions	4° C, store in dark, DO NOT FREEZE!
RRID	AB_2574145

Applications	Tested Dilution	Publications
Immunohistochemistry (IHC)	-	2 Publications
Flow Cytometry (Flow)	5 μL (0.125 μg)/test	7 Publications

Product Specific Information

Description: The dG9 antibody clone reacts with human perforin (pore-forming protein, pfp). Perforin is one of the cytolytic mediators present in the cytoplasmic granules of cytotoxic T lymphocytes (CTL) and natural killer cells (NK). Perforin is involved in the killing function by CTLs and NKs and has an important role in the immune response against tumors and virus infections.

Applications Reported: This dG9 (delta G9) antibody has been reported for use in intracellular staining followed by flow cytometric analysis.


Applications Tested: This dG9 (delta G9) antibody has been pre-titrated and tested by intracellular staining followed by flow cytometric analysis of normal human peripheral blood cells using the Intracellular Fixation & Permeabilization Buffer Set (Product # 88-8824-00) and protocol. Please refer to Best Protocols: Protocol A: Two step protocol for (cytoplasmic) intracellular proteins located under the Resource Tab online. This can be used at 5 μ L (0.125 μ g) per test. A test is defined as the amount (μ g) of antibody that will stain a cell sample in a final volume of 100 μ L. Cell number should be determined empirically but can range from 10^5 to 10^8 cells/test.

eFluor® 450 is an alternative to Pacific Blue®. eFluor® 450 emits at 445 nm and is excited with the Violet laser (405 nm). Please make sure that your instrument is capable of detecting this fluorochome.

Excitation: 405 nm; Emission: 445 nm; Laser: Violet Laser.

Filtration: 0.2 µm post-manufacturing filtered.

Product Images For Perforin Monoclonal Antibody (dG9 (delta G9)), eFluor™ 450, eBioscience™



Perforin Antibody (48-9994-42) in Flow

Staining of normal human peripheral blood cells with Anti-Human CD56 (NCAM) APC (Product # 17-0567-42) and Mouse IgG2b K Isotype Control eFluor® 450 (Product # 48-4732-82) (left) or Anti-Human Perforin eFluor® 450 (right) using the Intracellular Fixation and Permeabilization Buffer Set (Product # 88-8824-00) and protocol. Cells in the lymphocyte gate were used for analysis.

View more figures on thermofisher.cn

□ 9 References

Immunohistochemistry (2)

Nature communications

Single-cell transcriptomics links malignant T cells to the tumor immune landscape in cutaneous T cell lymphoma.

Year 2022

"Published figure using Perforin monoclonal antibody (Product # 48-9994-42) in Immunohistochemistry"

Authors: Liu X,Jin S,Hu S,Li R,Pan H,Liu Y,Lai P,Xu D,Sun J,Liu Z,Gao Y,Zhao Y,Liu F,Xiao Y,Li Y,Wen Y,Chen Z,Xu B,Lin Y,Ran M,Li Q,Yang S,Li H,Tu P,Haniffa M,Teichmann SA,Bai F,Wang Y

Science translational medicine

Lipoxin A4 regulates natural killer cell and type 2 innate lymphoid cell activation in asthma.

"Published figure using Perforin monoclonal antibody (Product # 48-9994-42) in Immunofluorescence"

Authors: Barnig C,Cernadas M,Dutile S,Liu X,Perrella MA,Kazani S,Wechsler ME,Israel E,Levy BD

Year 2013

Flow Cytometry (7)

Cancer immunology research

CD8+ T cell-Dependent Remodeling of the Tumor Microenvironment Overcomes Chemoresistance.

"48-9994-42 was used in Flow cytometry/Cell sorting to identify a subset of CD161-overexpressing CD8+ T cells enriched in chemoresistant tumors. CD161 engagement repressed the calcium influx and cytolytic capacity of CD8+ T cells through acid sphingomyelinase activation and ceramide generation."

Authors: Lao L, Zeng W, Huang P, Chen H, Jia Z, Wang P, Huang D, Chen J, Nie Y, Yang L, Wu W, Liu J

Year 2023

Species Human

JCI insight

Rapamycin improves Graves' orbitopathy by suppressing CD4+ cytotoxic T lymphocytes.

"Published figure using Perforin monoclonal antibody (Product # 48-9994-42) in Flow Cytometry"

Authors: Zhang M,Chong KK,Chen ZY,Guo H,Liu YF,Kang YY,Li YJ,Shi TT,Lai KK,He MQ,Ye K,Kahaly GJ,Shi BY, Wang Y

Year 2023

View more Flow references on thermofisher.cn

More applications with references on thermofisher.cn

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization, Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications requised by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless of service is imitted to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPRESS OR IMPLED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT.

BUYER'S EXCLUSIVE REMEDY FOR NON-CORNOR/MINO PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED. ACREMENT OF OR REFUND FOR THE NON-CONFORMINO PRODUCTS, SI AT SELECTIVES, SOLE OPTION. THERE IS NO ALGORITHM THE WARRANTY PERIOD IS LIMITED. AND THE PRODUCTS AS THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, or any type of consumption to human or animal to have the product is in