

Glucocorticoid Receptor Monoclonal Antibody (BuGR2)

Product Details

Size	100 µg
Species Reactivity	Human, Mouse, Sheep, Rabbit, Rat, Yeast, Guinea pig
Published Species	Rabbit, Rat, Yeast, Non-human primate, Sheep, Mouse, Human, Xenopus
Host/Isotype	Mouse / IgG2a
Class	Monoclonal
Type	Antibody
Clone	BuGR2
Conjugate	Unconjugated
Immunogen	Partially purified rat GR.
Form	Lyophilized
Concentration	1 mg/mL
Purification	Protein A
Storage buffer	PBS, pH 7.2
Contains	0.05% sodium azide
Storage conditions	-20° C, Avoid Freeze/Thaw Cycles
RRID	AB_325427

Applications	Tested Dilution	Publications
Western Blot (WB)	5 µg/mL	72 Publications
Immunohistochemistry (IHC)	-	9 Publications
Immunohistochemistry (Paraffin) (IHC (P))	5 µg/mL	-
Immunohistochemistry (PFA fixed) (IHC (PFA))	1:500	-
Immunohistochemistry (Frozen) (IHC (F))	-	1 Publication
Immunocytochemistry (ICC/IF)	Assay-dependent	13 Publications
Flow Cytometry (Flow)	1-2 µg/test	4 Publications
ELISA (ELISA)	-	2 Publications
Immunoprecipitation (IP)	Assay-dependent	22 Publications
ChIP assay (ChIP)	-	19 Publications
Gel Shift (GS)	Assay-dependent	5 Publications
Miscellaneous PubMed (Misc)	-	1 Publication

Product Specific Information

MA1-510 detects glucocorticoid receptor (GR) from human, mouse, rat, guinea pig, rabbit, sheep and yeast samples. This antibody does not react with primate, avian or amphibian GR.

MA1-510 has been successfully used in Western blot, immunofluorescence, immunocytochemistry, flow cytometry, immunohistochemistry, immunoprecipitation, and gel shift procedures. By Western blot, this antibody detects a 97 kDa protein representing GR in L929 cell extract. Immunocytochemical staining of GR in L929 cells with MA1-510 results in staining of both the cytoplasm and nucleus, even in the presence of hormone. Using enzymatic digestion analysis, MA1-510 reacts with the

undigested 97 kDa GR, a 17 kDa DNA-binding trypsin fragment, and a 45 kDa steroid- and DNA-binding chymotrypsin fragment.

The MA1-510 immunogen is partially purified rat GR.

Reconstitute with 100 µL distilled water.

Product Images For Glucocorticoid Receptor Monoclonal Antibody (BuGR2)

Glucocorticoid Receptor Antibody (MA1-510) in ICC/IF

Immunofluorescent analysis of Glucocorticoid Receptor using Glucocorticoid Receptor Monoclonal Antibody (BuGR2) (Product # MA1-510) shows staining in A549 Cells. Glucocorticoid Receptor (green), F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue) is shown. Cells were grown on chamber slides and fixed with formaldehyde prior to staining. Cells were probed without (control) or with an antibody recognizing Glucocorticoid Receptor (Product # MA1-510) at a dilution of 1:100 over night at 4 °C, washed with PBS and incubated with a DyLight-488 conjugated secondary antibody (Product # 35552 for GAR, Product # 35503 for GAM). Images were taken at 60X magnification.

Glucocorticoid Receptor Antibody (MA1-510) in ICC/IF

Immunofluorescent analysis of Glucocorticoid Receptor using Glucocorticoid Receptor Monoclonal Antibody (BuGR2) (Product # MA1-510) shows staining in U251 Cells. Glucocorticoid Receptor (green), F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue) is shown. Cells were grown on chamber slides and fixed with formaldehyde prior to staining. Cells were probed without (control) or with an antibody recognizing Glucocorticoid Receptor (Product # MA1-510) at a dilution of 1:100 over night at 4 °C, washed with PBS and incubated with a DyLight-488 conjugated secondary antibody (Product # 35552 for GAR, Product # 35503 for GAM). Images were taken at 60X magnification.

Glucocorticoid Receptor Antibody (MA1-510) in ICC/IF

Immunofluorescent analysis of Glucocorticoid Receptor using Glucocorticoid Receptor Monoclonal Antibody (BuGR2) (Product # MA1-510) shows staining in Hela Cells. Glucocorticoid Receptor (green), F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue) is shown. Cells were grown on chamber slides and fixed with formaldehyde prior to staining. Cells were probed without (control) or with an antibody recognizing Glucocorticoid Receptor (Product # MA1-510) at a dilution of 1:100 over night at 4 °C, washed with PBS and incubated with a DyLight-488 conjugated secondary antibody (Product # 35552 for GAR, Product # 35503 for GAM). Images were taken at 60X magnification.

View more figures on thermofisher.cn

Western Blot (72)

<p>Frontiers in molecular neuroscience</p> <p>Prolonged maternal exposure to glucocorticoids alters selenoprotein expression in the developing brain.</p> <p>"Published figure using Glucocorticoid Receptor monoclonal antibody (Product # MA1-510) in Western Blot"</p> <p>Authors: Toh P,Seale LA,Berry MJ,Torres DJ</p>	<p>Year</p> <p>2024</p> <p>Species</p> <p>Mouse</p> <p>Dilution</p> <p>1:1,000</p>
<p>Cells</p> <p>Aurora Kinase A Regulates Cell Transitions in Glucocorticoid-Induced Bone Loss.</p> <p>"Published figure using Glucocorticoid Receptor monoclonal antibody (Product # MA1-510) in Western Blot"</p> <p>Authors: Qiao X,Yang Y,Zhao Y,Wu X,Zhang L,Cai X, Ji J,Boström KI,Yao Y</p>	<p>Year</p> <p>2023</p>

[View more WB references on thermofisher.cn](#)

Immunohistochemistry (9)

<p>iScience</p> <p>Loss of central mineralocorticoid or glucocorticoid receptors impacts auditory nerve processing in the cochlea.</p> <p>"MA1-510 was used in Immunohistochemistry to imply that stress hormone-dependent functions of central MR/GR contribute to "precognitive" sound processing in the cochlea."</p> <p>Authors: Marchetta P,Eckert P,Lukowski R,Ruth P,Singer W,Rüttiger L,Knipper M</p>	<p>Year</p> <p>2022</p> <p>Species</p> <p>Mouse</p> <p>Dilution</p> <p>1:500</p>
<p>Reproductive sciences (Thousand Oaks, Calif.)</p> <p>Effects of maternal dexamethasone treatment early in pregnancy on glucocorticoid receptors in the ovine placenta.</p> <p>"Published figure using Glucocorticoid Receptor monoclonal antibody (Product # MA1-510) in Immunohistochemistry"</p> <p>Authors: Shang H,Meng W,Sloboda DM,Li S,Ehrlich L,Plagemann A,Dudenhause JW, Henrich W,Newnham JP,Challis JR,Braun T</p>	<p>Year</p> <p>2015</p> <p>Species</p> <p>Sheep</p> <p>Dilution</p> <p>1:50</p>

[View more IHC references on thermofisher.cn](#)

More applications with references on thermofisher.cn

- IHC (F) (1)
- ICC/IF (13)
- Flow (4)
- ELISA (2)
- IP (22)
- ChIP (19)
- GS (5)
- Misc (1)

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPRESS OR IMPLIED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT. BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED TO REPAIR, REPLACEMENT OF OR REFUND FOR THE NON-CONFORMING PRODUCT(S) AT SELLER'S SOLE OPTION. THERE IS NO OBLIGATION TO REPAIR, REPLACE OR REFUND FOR PRODUCTS AS THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses, or any type of consumption by or application to human or animals.