



# **IDO1** Recombinant Rabbit Monoclonal Antibody (BLR031F)


<b>Product Details</b>	
Size	100 μL
Species Reactivity	Human
Host/Isotype	Rabbit / IgG
Expression system	HEK293 cells
Class	Recombinant Monoclonal
Туре	Antibody
Clone	BLR031F
Conjugate	Unconjugated
Immunogen	Residues 1-403 (Full Length)
Form	Liquid
Concentration	50 μg/mL
Purification	purified
Storage buffer	BBS, pH 8.2, with 0.1% BSA
Contains	0.09% sodium azide
Storage conditions	2-8°C
RRID	AB_2926387

Applications	Tested Dilution	Publications
Western Blot (WB)	1:1,000	-
Immunohistochemistry (Paraffin) (IHC (P))	1:100-1:500	-
Immunocytochemistry (ICC/IF)	1:100-1:500	-

# **Product Specific Information**


Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE cell sections.

# Product Images For IDO1 Recombinant Rabbit Monoclonal Antibody (BLR031F)




#### IDO1 Antibody (MA5-44257) in WB

Western Blot of IDO1 in whole cell lysate (50 µg) from HeLa treated with IFN-gamma (+) or mock treated (-), HEK293T, HDLM-2, Hep-G2, IDO2 over-expressing HEK, untransfected HEK, and IDO1 over-expressing HEK cells. Samples were treated with NETN lysis buffer, incubated with IDO1 monoclonal antibody (Product # MA5-44258) at a dilution of 1:1000, followed by HRP-conjugated goat anti-rabbit IgG (exposure time of 30 seconds) Lower Panel: Rabbit anti-Vinculin.



## IDO1 Antibody (MA5-44257) in ICC/IF

Immunocytochemistry analysis of IDO1 in A-549 cells, untreated (left) or treated with Interferon-gamma (right). Samples were incubated with IDO1 monoclonal antibody (Product # MA5-44258) at a dilution of 1:250 followed by HRP-conjugated goat anti-rabbit IgG and DAB.



## IDO1 Antibody (MA5-44257) in IHC (P)

Immunohistochemistry analysis of IDO1 in paraffin-embedded skin carcinoma. Samples were incubated with IDO1 monoclonal antibody (Product # MA5-44258) at a dilution of 1:250, HRP-conjugated goat anti-rabbit IgG, Opal substrate and DAPI.

## View more figures on thermofisher.cn

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production specifications and/or accompanying package inserts ("Documentation"), No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is imited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anynow other than the Buyer. Any mode of sample invitable to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample invitable. The product is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPRESS OR IMPLED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES FOR REFORM PARTICULAR PURPOSE, OR NON INFRINGEMENT.

BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED TO REPAIR, REPLACEMENT OF OR REFUND FOR THE NON-CONFORMING PRODUCTS, AT SELLER'S SOLD FOR INTERED AND CONFORMING PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, or any type of consumption to human or animal to how themselved uses, or any type of consumption to human or animal to how the animal transfer of consumption to human or animal transfer of the purpose, including witho