

TRPC7 Monoclonal Antibody (N64A/36), APC


Product Details	
Size	100 μg
Species Reactivity	Human, Mouse, Rat
Host/Isotype	Mouse / IgG1
Class	Monoclonal
Туре	Antibody
Clone	N64A/36
Conjugate	APC
Excitation/Emission Max	651/660 nm
Immunogen	Synthetic peptide amino acids 845-862 of human TRPC7
Form	Liquid
Concentration	1 mg/mL
Purification	Protein G
Storage buffer	47.35mM phosphate/2.15mM MES, pH 7.4, with 1.033mM EDTA
Contains	no preservative
Storage conditions	4° C

Applications	Tested Dilution	Publications
Western Blot (WB)	1:1,000	-
Immunohistochemistry (Paraffin) (IHC (P))	1:100	-
Immunocytochemistry (ICC/IF)	1:100	-
Immunoprecipitation (IP)	Assay-dependent	-
Antibody Microarray (AM)	Assay-dependent	-

Product Specific Information


1 μg/mL of MA5-45473 was sufficient for detection of TrpC7 in 10 μg of rat brain lysate by colorimetric immunoblot analysis using goat anti-mouse IgG:HRP as the secondary antibody.| Detects approximately 100kDa.

Product Images For TRPC7 Monoclonal Antibody (N64A/36), APC




TRPC7 Antibody (MA5-45473) in ICC/IF

Immunocytochemistry/Immunofluorescence analysis using human neuroblastoma cells. Fixation involved 4% PFA for 15 min. Samples were incubated with TrpC7 monoclonal antibody (Product # MA5-45473) at 1:100 for overnight at 4°C with slow rocking, followed by AlexaFluor 488 at 1:1,000 for 1 hour at RT. Counterstain used was Phalloidin-iFluor 647 (red) F-Actin stain; Hoechst (blue) nuclear stain at 1:800, 1.6mM for 20 min at RT. (A) Hoechst (blue) nuclear stain. (B) Phalloidin-iFluor 647 (red) F-Actin stain. (C) TrpC7 antibody (D) Composite.



TRPC7 Antibody (MA5-45473) in ICC/IF

Immunocytochemistry/Immunofluorescence analysis using HaCaT cells. Fixation involved Cold 100% methanol for 10 minutes at -20°C. Samples were incubated with TrpC7 monoclonal antibody (Product # MA5-45473) at 1:100 for 1 hour at RT, followed by FITC Goat Anti-Mouse (green) at 1:50 for 1 hour at RT. Localization: Nuclear staining.



TRPC7 Antibody (MA5-45473) in IHC (P)

Immunohistochemistry analysis using backskin. Fixation involved Bouins Fixative and paraffin-embedded. Samples were incubated with TrpC7 monoclonal antibody (Product # MA5-45473) at 1:100 for 1 hour at RT, followed by FITC Goat Anti-Mouse (green) at 1:50 for 1 hour at RT. Localization: Everything.

View more figures on thermofisher.cn

For Research Use Only, Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production of specifications and/or accompanying package inserts ("Documentation"), No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is important to the product is used to any properly trained individuals. Unless otherwise stated in the Documentation, this warranty does not extend to anypone other than the Buyer. Any mode of a sample increase in merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample increase. NO OTHER WARRANTES, EXPRESS OR IMPLED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT.

BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED. A CREATED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT.

BUYER'S EXCLUSIVE EXEMBEDY FOR NON-CONFORMING PRODUCTS SOLIC STATE AND PRODUCTS OR THE NON-CONFORMING PRODUCTS OR THE NON-CONFORMING PRODUCTS, AS THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, or vivo or invivo therapeutic uses, or any ty