



Phospho-Ezrin (Tyr478) Polyclonal Antibody


Product Details		
Size	100 μL	
Species Reactivity	Human, Mouse, Rat	
Host/Isotype	Rabbit / IgG	
Class	Polyclonal	
Туре	Antibody	
Conjugate	Unconjugated	
Immunogen	A synthesized peptide derived from human EZR(Accession P15311), corresponding to amino acid residues around phosphorylated Tyr478.	
Form	Liquid	
Concentration	1 mg/mL	
Purification	sequential chromatography	
Storage buffer	PBS, pH 7.4, with 50% glycerol	
Contains	0.02% sodium azide	
Storage conditions	-20°C	
RRID	AB_2816273	

Applications	Tested Dilution	Publications
Western Blot (WB)	1:500-1:2,000	-
Immunohistochemistry (Paraffin) (IHC (P))	1:100-1:500	-
Immunocytochemistry (ICC/IF)	1:100-1:500	-

Product Specific Information


Antibody detects endogenous levels of Ezrin only when phosphorylated at Tyrosine 478.

Product Images For Phospho-Ezrin (Tyr478) Polyclonal Antibody




Phospho-Ezrin (Tyr478) Antibody (PA5-104800) in ICC/IF

Immunofluorescent analysis of Phospho-Ezrin (Tyr478) in HeLa cells(4h of LPS treatment). Samples were fixed with paraformaldehyde, permeabilized with 0.1% Triton X-100, blocked with 10% serum (45 min at 25°C), incubated with mouse anti-beta tubulin and Phospho-Ezrin (Tyr478) polyclonal antibody (Product # PA5-104800) using a dilution of 1:200 (1 hr, 37°C), and followed by goat anti-rabbit IgG Alexa Fluor 594 (red) and goat anti-mouse IgG Alexa Fluor 488 (green).



Phospho-Ezrin (Tyr478) Antibody (PA5-104800) in IHC (P)

Immunohistochemistry analysis of paraffin-embedded Phospho-Ezrin (Tyr478) in rat intestinal tissue sections. Antigen retrieval was performed using citrate buffer. Samples were blocked with blocking buffer (1.5 hr, 22°C), incubated with Phospho-Ezrin (Tyr478) polyclonal antibody (Product # PA5-104800) using a dilution of 1:200 (1.5 hr, 22°C), followed by HRP conjugated goat anti-rabbit.



Phospho-Ezrin (Tyr478) Antibody (PA5-104800) in IHC (P)

Immunohistochemistry analysis of paraffin-embedded Phospho-Ezrin (Tyr478) in rat intestinal tissue sections. Antigen retrieval was performed using citrate buffer. Samples were blocked with blocking buffer (1.5 hr, 22°C), incubated with Phospho-Ezrin (Tyr478) polyclonal antibody (Product # PA5-104800) using a dilution of 1:200 (1.5 hr, 22°C), followed by HRP conjugated goat anti-rabbit.

View more figures on thermofisher.cn

For Research Use Only, Not for use in diagnostic procedures. Not for resale without express authorization, Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless of services the use of subject to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPRESS OR IMPLED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT.

BUYER'S EXCLUSIVE REMEDY FOR NON-CORNORMOR PRODUCTS DURING THE WARRANTY PERFIOI DIS LIMITED. A REFLION FOR THE NON-CONFORMING PRODUCTS AS THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying is.