



## alpha-Fetoprotein Monoclonal Antibody (1E8), Biotin, eBioscience™

<b>Product Details</b>	
Size	100 μg
Species Reactivity	Human
Host/Isotype	Mouse / IgG1, kappa
Recommended Isotype Control	Mouse IgG1 kappa Isotype Control (P3.6.2.8.1), Biotin, eBioscience™
Class	Monoclonal
Туре	Antibody
Clone	1E8
Conjugate	Biotin
Form	Liquid
Concentration	0.5 mg/mL
Purification	Affinity chromatography
Storage buffer	PBS, pH 7.2
Contains	0.09% sodium azide
Storage conditions	4° C, store in dark, DO NOT FREEZE!
RRID	AB_2572812

Applications	Tested Dilution	Publications
Immunohistochemistry (Paraffin) (IHC (P))	20 μg/mL	-
Immunocytochemistry (ICC/IF)	Assay-Dependent	-

## **Product Specific Information**


Description: This 1E8 monoclonal antibody reacts with human alpha-fetoprotein (AFP). This 70-kDa secretory protein is a member of the albumin gene family. Synthesized by the yolk sac and fetal liver during embryogenesis, AFP protein levels are highest in fetal serum. After birth, serum AFP levels decrease dramatically. In fact, AFP is nearly undetectable in normal adult serum. However, hepatocellular carcinoma and germ cell teratoblastoma, as well as liver regeneration, viral hepatitis, and cirrhosis, leads to elevated AFP serum levels in adults. As such, detection of this protein is frequently used as a diagnostic marker for these conditions.


Applications Reported: This 1E8 antibody has been reported for use in microscopy, immunohistochemical staining, and immunocytochemistry.

Applications Tested: This 1E8 antibody has been tested by immunohistochemistry on formalin-fixed paraffin embedded tissue using high pH antigen retrieval at 20  $\mu$ g/mL. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

Filtration: 0.2 µm post-manufacturing filtered.

## Product Images For alpha-Fetoprotein Monoclonal Antibody (1E8), Biotin, eBioscience™





## alpha-Fetoprotein Antibody (13-9499-82) in IHC (P)

Immunohistochemistry of formalin-fixed paraffin embedded human liver stained with 20  $\mu$ g/mL Mouse IgG1 K Isotype Control Biotin (left) or 20  $\mu$ g/mL Anti-Human alpha-Fetoprotein Biotin (right) followed by streptavidin HRP and DAB visualization. Nuclei are counterstained with hematoxylin.

For Research Use Only, Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. No OTHER WARRANTIES, EXPERESS OR IMPLECT LIMITATION, IMPLIED WARRANTIES, EXPERESS OR IMPLECT. IN IMPLIED WARRANTIES, EXPERESS OR IMPLECT. IN IMPLICATION, IMPLIED WARRANTIES, EXPERESS OR IMPLECT. IN IMPLICATION, IMPLIED WARRANTIES, EXPERESS OR IMPLECT. IN IMPLICATION, IMPL