CD62L (L-Selectin) Monoclonal Antibody (MEL-14), PerCP-Cyanine5.5, eBioscience™

| Product Details | |
|--------------------------------|---|
| Size | 100 µg |
| Species Reactivity | Mouse |
| Published Species | Rat, Mouse |
| Host/Isotype | Rat / IgG2a, kappa |
| Recommended Isotype Control | Rat IgG2a kappa Isotype Control (eBR2a), PerCP-Cyanine5.5, eBioscience™ |
| Class | Monoclonal |
| Туре | Antibody |
| Clone | MEL-14 |
| Conjugate | PerCP-Cyanine5.5 |
| Excitation/Emission Max | 489/679 nm |
| Form | Liquid |
| Concentration | 0.2 mg/mL |
| Purification | Affinity chromatography |
| Storage buffer | PBS, pH 7.2 |
| Contains | 0.09% sodium azide |
| Storage conditions | 4° C, store in dark, DO NOT FREEZE! |
| RRID | AB_996667 |

| Applications | Tested Dilution | Publications |
|----------------------------|-----------------|-----------------|
| Immunohistochemistry (IHC) | - | 1 Publication |
| Flow Cytometry (Flow) | 0.125 μg/test | 62 Publications |

Product Specific Information

Description: The MEL-14 monoclonal antibody reacts with mouse CD62L, a 76 kDa member of the selectin family. CD62L is expressed by neutrophils, monocytes, and subsets of T, B, and NK cells and binds a number of glycosylated, fucosylated, sulfated sialylated glycoproteins including CD34, glycam-1 and MAdCam-1. These interactions mediate rolling of lymphocytes on activated endothelium at the sites of inflammation and homing of cells to the high endothelial venules (HEV) of peripheral lymphoid tissues.

Applications Reported: This MEL-14 antibody has been reported for use in flow cytometric analysis.


Applications Tested: This MEL-14 antibody has been tested by flow cytometric analysis of mouse splenocytes. This can be used at less than or equal to 0.125 μ g per test. A test is defined as the amount (μ g) of antibody that will stain a cell sample in a final volume of 100 μ L. Cell number should be determined empirically but can range from 10^5 to 10^8 cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

Excitation: 488 nm; Emission: 695 nm; Laser: Blue Laser.

Filtration: 0.2 µm post-manufacturing filtered.

1

Product Images For CD62L (L-Selectin) Monoclonal Antibody (MEL-14), PerCP-Cyanine5.5, eBioscience™



CD62L (L-Selectin) Antibody (45-0621-82) in Flow

Staining of BALB/c splenocytes with 0.06 µg of Rat IgG2a K Isotype Control PerCP-Cyanine5-5 (Product # 45-4321-80) (open histogram) or 0.06 µg of Anti-Mouse CD62L (L-Selectin) PerCP-Cyanine5-5 (filled histogram). Total viable cells were used for analysis.

View more figures on thermofisher.cn

| A 63 References | |
|--|---|
| mmunohistochemistry (1) | |
| Nature communications Intratumoral immunotherapy using platelet-cloaked nanoparticles enhances antitumor immunity in solid tumors. "45-0621-82 was used in Immunohistochemistry to highlight the promise of locally delivering immunostimulatory payloads using biomimetic nanocarriers, which possess advantages such as enhanced biocompatibility and natural targeting affinities." Authors: Bahmani B,Gong H,Luk BT,Haushalter KJ,DeTeresa E,Previti M,Zhou J,Gao W,Bui JD,Zhang L,Fang RH, Zhang J | Year 2021 Species Mouse Dilution 1:400 |
| low Cytometry (62) | |
| Bioactive materials | Year 2023 |
| A platinum@polymer-catechol nanobraker enables radio- immunotherapy for crippling melanoma tumorigenesis, angiogenesis, and radioresistance. | 2020 |
| "Published figure using CD62L (L-Selectin) monoclonal antibody (Product # 45-0621-82) in Flow Cytometry" | |

| NPJ vaccines Amphiphile-CpG vaccination induces potent lymph node activation and COVID-19 immunity in mice and non-human primates. | Year 2022 |
|--|---------------------|
| "Published figure using CD62L (L-Selectin) monoclonal antibody (Product # 45-0621-82) in Flow Cytometry" | |
| Authors: Seenappa LM,Jakubowski A,Steinbuck MP,Palmer E,Haqq CM,Carter C,Fontenot J,Villinger F,McNeil LK, DeMuth PC | |

View more Flow references on thermofisher.cn

More applications with references on thermofisher.cn

Authors: Li W, Yan J, Tian H, Li B, Wang G, Sang W, Zhang Z, Zhang X, Dai Y

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Those warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not

2

represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPRESS OR IMPLED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT. BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED TO REPAIR, REPLACEMENT OF OR REFUND FOR THE NON-CONFORMING PRODUCT(S) AT SELLER'S SOLE OPTION. THERE IS NO OBLIGATION TO REPAIR, REPLACE OR REFUND FOR PRODUCTS AS THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. UNless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses, or any type of consumption by or application to human or animals.