Performance guarenteed

# LIN7C Polyclonal Antibody


## **Product Details**

Size	100 µg
Species Reactivity	Dog, Human, Mouse, Non-human primate, Rat
Published Species	Mouse, Human
Host/Isotype	Rabbit / IgG
Class	Polyclonal
Туре	Antibody
Conjugate	Unconjugated
Immunogen	Synthetic peptide derived from the C-terminus of the rat Velis-3 (MALS-3) protein.
Form	Liquid
Concentration	0.25 mg/mL
Purification	Antigen affinity chromatography
Storage buffer	PBS, pH 7.4
Contains	0.1% sodium azide
Storage conditions	-20° C, Avoid Freeze/Thaw Cycles
RRID	AB_2533910

Applications	Tested Dilution	Publications
Western Blot (WB)	2 µg/mL	5 Publications
Immunohistochemistry (IHC)	-	2 Publications
Immunohistochemistry (Frozen) (IHC (F))	1.0 µg/mL	3 Publications
Immunocytochemistry (ICC/IF)	2 μg/mL	5 Publications
ELISA (ELISA)	0.1-1.0 μg/mL	-
Immunoprecipitation (IP)	1 µg	1 Publication
Miscellaneous PubMed (Misc)	-	1 Publication

1


# Product Images For LIN7C Polyclonal Antibody



## LIN7C Antibody (51-5600) in WB

Western blot analysis was performed on whole cell extracts (30 µg lysate) of Neuro-2a (Lane 1), SH-SY5Y (Lane 2), U-87 MG (Lane 3), HEK 293 (Lane 4), Caki-1 (Lane 5), COS-7 (Lane 6), and MDCK (Lane 7). The blots were probed with Rabbit Anti-Veli-3 Polyclonal Antibody (Product # 51-5600, 2 µg/mL) and detected by chemiluminescence using Goat anti-Rabbit IgG (Heavy Chain) Superclonal<sup>™</sup> Secondary Antibody, HRP conjugate (Product # A27036, 0.25 µg /mL, 1:4000 dilution). A 22 kDa band corresponding to LIN7C (Veli-3) was observed across the cell lines tested. Known quantity of protein samples were electrophoresed using Novex® NuPAGE® 4-12 % Bis-Tris gel (Product # NP0321BOX), XCell SureLock™ Electrophoresis System (Product # El0002) and Novex® Sharp Pre-Stained Protein Standard (Product # LC5800). Resolved proteins were then transferred onto a nitrocellulose membrane with iBlot® 2 Dry Blotting System (Product # IB21001). The membrane was probed with the relevant primary and secondary Antibody following blocking with 5 % skimmed milk. Chemiluminescent detection was performed using Pierce™ ECL Western Blotting Substrate (Product # 32106).

#### LIN7C Antibody (51-5600) in ICC/IF



Immunofluorescence analysis of VELI-3 was performed using 70% confluent log phase RSC96 cells. The cells were fixed with 4% paraformaldehyde for 10 minutes, permeabilized with 0.1% Triton<sup>™</sup> X-100 for 10 minutes, and blocked with 1% BSA for 1 hour at room temperature. The cells were labeled with LIN7C Rabbit Polyclonal Antibody (Product # 51-5600) at 2µg/mL in 0.1% BSA and incubated for 3 hours at room temperature and then labeled with Goat anti-Rabbit IgG (Heavy Chain) Superclonal<sup>™</sup> Secondary Antibody, Alexa Fluor® 488 conjugate (Product # A27034) at a dilution of 1:2000 for 45 minutes at room temperature (Panel a: green). Nuclei (Panel b: blue) were stained with SlowFade® Gold Antifade Mountant with DAPI (Product # S36938). F-actin (Panel c: red) was stained with Alexa Fluor® 555 Rhodamine Phalloidin (Product # R415, 1:300). Panel d represents the merged image showing membranous localization. Panel e shows the no primary antibody control. The images were captured at 60X magnification.

View more figures on thermofisher.cn

2

## **□** 17 References

## Western Blot (5)

Journal of psychiatry & neuroscience : JPN Calcium/calmodulin-dependent serine protein kinase (CASK), a protein implicated in mental retardation and autism-spectrum disorders, interacts with T-Brain-1 (TBR1) to control extinction of associative memory in male mice.	Year 2017 <b>Species</b> Mouse
"51-5600 was used in Western Blotting to establish a mouse model to study how calcium/calmodulin-dependent serine protein kinase regulates mental ability."	
Authors: Huang TN,Hsueh YP	
The American journal of pathology	<b>Year</b> 2013

miR-199a-5p regulates urothelial permeability and may play a role in bladder pain syndrome.

"51-5600 was used in western blot to show that miR-99a-5p regulates intercellular junctions in the bladder." Authors: Monastyrskaya K,Sánchez-Freire V,Hashemi Gheinani A,Klumpp DJ,Babiychuk EB,Draeger A,Burkhard FC

View more WB references on thermofisher.cn

Species Human

# Immunohistochemistry (2)

PloS one	Year
Testing for a gap junction-mediated bystander effect in retinitis	2013
pigmentosa: secondary cone death is not altered by deletion of	
connexin36 from cones.	Dilution 1:1000
51-5600 was used in immunohistochemistry to test if disruption of rod-cone coupling prevents secondary cone death and reduces the spread of degeneration"	
Authors: Kranz K,Paquet-Durand F,Weiler R,Janssen-Bienhold U,Dedek K	
Nolecular biology of the cell	Year
	<b>Year</b> 2009
Molecular biology of the cell CASK deletion in intestinal epithelia causes mislocalization of LIN7C and the DLG1/Scrib polarity complex without affecting cell polarity.	
CASK deletion in intestinal epithelia causes mislocalization of LIN7C	

## More applications with references on thermofisher.cn

IHC (F) (3) ICC/IF (5) IP (1) Misc (1)

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warrant y provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warrant y is used or sample furnished to sample furnished sample sample for the sample sample for the produc