


4EBP1 Monoclonal Antibody (554R16)

Product Details	
Size	100 µg
Species Reactivity	Human, Mouse, Rat
Published Species	Rat, Human
Host/Isotype	Mouse / IgG1, kappa
Class	Monoclonal
Type	Antibody
Clone	554R16
Conjugate	Unconjugated
Immunogen	Recombinant human 4E-BP1 protein expressed in E. coli.
Form	Liquid
Concentration	0.5 mg/mL
Purification	purified
Storage buffer	PBS, pH 7.2, with 1% BSA
Contains	0.1% sodium azide
Storage conditions	-20°C
RRID	AB_2536336

Applications	Tested Dilution	Publications
Western Blot (WB)	1:500	4 Publications
Immunocytochemistry (ICC/IF)	1:250	1 Publication
Miscellaneous PubMed (Misc)	-	1 Publication

Product Images For 4EBP1 Monoclonal Antibody (554R16)




4EBP1 Antibody (AHO1382)

Antibody specificity was demonstrated by siRNA mediated knockdown of target protein. A-431 cells were transfected with 4EBP1 siRNA and decrease in signal was observed in Western Blot using Anti-4EBP1 Monoclonal Antibody (Product # AHO1382). {KD}


4EBP1 Antibody (AHO1382) in WB

Western blot analysis was performed on whole cell extracts of A-431 (Lane 1), HEK-293 (Lane 2), HeLa (Lane 3), Hep G2 (Lane 4) and U-87 MG (Lane 5). The blot was probed with Anti-4EBP1 antibody (Product # AHO1382, 1:500 dilution) and detected by chemiluminescence using Goat anti-Mouse IgG (H+L) Superclonal™ Secondary Antibody, HRP conjugate (Product # A28177, 0.25 µg/mL, 1:4000 dilution). A 17 kDa band corresponding to 4EBP1 was observed across the cell lines tested.



4EBP1 Antibody (AHO1382) in ICC/IF

Immunofluorescence analysis of 4EBP1 was performed using 70% confluent log phase A-431 cells. The cells were fixed with 4% paraformaldehyde for 10 minutes, permeabilized with 0.1% Triton™ X-100 for 10 minutes, and blocked with 1% BSA for 1 hour at room temperature. The cells were labeled with 4EBP1 Monoclonal Antibody (Product # AHO1382) at 1:250 dilution in 0.1% BSA and incubated overnight at 4 degree and then labeled with Goat anti-Mouse IgG (H+L) Superclonal™ Secondary Antibody, Alexa Fluor® 488 conjugate (Product # A28175) at a dilution of 1:2000 for 45 minutes at room temperature (Panel a: green). Nuclei (Panel b: blue) were stained with SlowFade® Gold Antifade Mountant with DAPI (Product # S36938). F-actin (Panel c: red) was stained with Rhodamine Phalloidin (Product # R415, 1:300). Panel d represents the merged image showing nuclear and cytoplasmic localization. Panel e represents control cells with no primary antibody to assess background. The images were captured at 60X magnification.



View more figures on thermofisher.cn

6 References

Western Blot (4)

Cell death and differentiation

Inhibition of the autophagic protein ULK1 attenuates axonal degeneration in vitro and in vivo, enhances translation, and modulates splicing.

"AHO1382 was used in Western Blotting to study the effect of ULK1 inhibition in different models of lesion-induced axonal degeneration in vitro and in vivo."

Authors: Vahsen BF,Ribas VT,Sundermeyer J,Boecker A,Dambeck V,Lenz C,Shomroni O,Caldi Gomes L,Tatenhorst L, Barski E,Roser AE,Michel U,Urlaub H,Salinas G,Bähr M,Koch JC,Lingor P

Year
2020

Species
Rat

Dilution
1:200

Journal of cellular and molecular medicine

Polycystin-1 affects cancer cell behaviour and interacts with mTOR and Jak signalling pathways in cancer cell lines.

"AHO1382 was used in Western Blotting to study the role of polycystin-1 in cancer biology."

Authors: Papavassiliou KA,Zoi I,Gargalionis AN,Koutsilieris M

Year
2019

Species
Human

Dilution
1:250

[View more WB references on thermofisher.cn](#)

Immunocytochemistry (1)

The Journal of biological chemistry

Butyrate-rich colonic microenvironment is a relevant selection factor for metabolically adapted tumor cells.

Authors: Serpa J,Caiado F,Carvalho T,Torre C,Gonçalves LG,Casalou C,Lamosa P,Rodrigues M,Zhu Z,Lam EW,Dias S

Year
2010

Species
Human

Miscellaneous PubMed (1)

Autophagy

Induction of genuine autophagy by cationic lipids in mammalian cells.

"AHO1382 was used in western blot to report that cationic lipids induced autophagy in mammalian cells"

Authors: Man N,Chen Y,Zheng F,Zhou W,Wen LP

Year
2010

Species
Human

More applications with references on thermofisher.cn

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPRESS OR IMPLIED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON-INFRINGEMENT. BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED TO REPAIR, REPLACEMENT OF OR REFUND FOR THE NON-CONFORMING PRODUCT(S) AT SELLER'S SOLE OPTION. THERE IS NO OBLIGATION TO REPAIR, REPLACE OR REFUND FOR PRODUCTS AS THE RESULT OF (i) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (ii) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (iii) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (iv) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses, or any type of consumption by or application to human or animals.