

CD11b/c Monoclonal Antibody (OX-42)

Product Details

Size	100 µg
Species Reactivity	Rat
Published Species	Rat
Host/Isotype	Mouse / IgG2a
Class	Monoclonal
Type	Antibody
Clone	OX-42
Conjugate	Unconjugated
Immunogen	Rat peritoneal macrophages.
Form	Liquid
Concentration	1 mg/mL
Purification	Protein A
Storage buffer	PBS, pH 7.2
Contains	0.09% sodium azide
Storage conditions	Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.
RRID	AB_927470

Applications	Tested Dilution	Publications
Immunohistochemistry (IHC)	-	3 Publications
Immunohistochemistry (Frozen) (IHC (F))	1:50-1:100	1 Publication
Immunohistochemistry - Free Floating (IHC (Free))	-	1 Publication
Immunocytochemistry (ICC/IF)	Assay-dependent	2 Publications
Flow Cytometry (Flow)	1:50-1:100	4 Publications
Immunoprecipitation (IP)	Assay-dependent	1 Publication
Miscellaneous PubMed (Misc)	-	12 Publications

Product Specific Information

For FACS analysis, use 10 µL of the suggested working dilution to label 1×10^6 cells in 100 µL.

Mouse anti Rat CD11b, clone OX-42, recognizes rat CD11b, also known as integrin alpha-M, the receptor for the iC3b component of complement. CD11b is a 1151 amino acid single pass type 1 transmembrane glycoprotein possessing a single vWFA domain and multiple FG-GAP repeats.

Mouse anti Rat CD11b, clone OX-42, recognizes rat CD11b, also known as integrin alpha-M, the receptor for the iC3b component of complement. CD11b is a 1151 amino acid single pass type 1 transmembrane glycoprotein possessing a single vWFA domain and multiple FG-GAP repeats.

Immunohistochemistry (3)

<p>International journal of molecular sciences</p> <p>Differential Expression of Neuroinflammatory mRNAs in the Rat Sciatic Nerve Following Chronic Constriction Injury and Pain-Relieving Nanoemulsion NSAID Delivery to Infiltrating Macrophages.</p> <p>"MA1-81606 was used in Immunohistochemistry to the prognostic significance of the LMR associated with RT in patients with WR-B-NHL and acknowledged the potential use of PD-1 antibody in RT-treated lymphomas."</p> <p>Authors: Stevens AM,Liu L,Bertovich D,Janjic JM,Pollock JA</p>	<p>Year 2019</p> <p>Dilution 1:100</p>
<p>Journal of neuroscience research</p> <p>Increased ceruloplasmin expression caused by infiltrated leukocytes, activated microglia, and astrocytes in injured female rat spinal cords.</p> <p>"MA1-81606 was used in Immunohistochemistry to show how quantitative analysis proved that infiltrated leukocytes, activated microglia/macrophages, and astrocytes should be the major sources of increased Cp."</p> <p>Authors: Wu Y,Shen L,Wang R,Tang J,Ding SQ,Wang SN,Guo XY,Hu JG,Lü HZ</p>	<p>Year 2018</p> <p>Species Rat</p> <p>Dilution 1:200</p>

[View more IHC references on thermofisher.cn](#)

Immunohistochemistry (Frozen) (1)

<p>Journal of neurotrauma</p> <p>Fluoxetine prevents oligodendrocyte cell death by inhibiting microglia activation after spinal cord injury.</p> <p>"MA1-81606 was used in immunohistochemistry - frozen section to demonstrate that fluoxetine alleviates oligodendrocyte cell death after spinal cord injury"</p> <p>Authors: Lee JY,Kang SR,Yune TY</p>	<p>Year 2015</p> <p>Dilution 1:200</p>
--	--

Immunohistochemistry - Free Floating (1)

<p>Neuropathology : official journal of the Japanese Society of Neuropathology</p> <p>A rat model of mild cognitive impairment associated with vascular factor.</p> <p>"MA1-81606 was used in immunohistochemistry - free floating to develop and characterize a rat model of vascular mild cognitive impairment"</p> <p>Authors: Zhang L,Wei WS,Li YJ,Wang Y</p>	<p>Year 2011</p> <p>Species Rat</p> <p>Dilution 1:50</p>
---	---

More applications with references on thermofisher.cn

ICC/IF (2) Flow (4) IP (1) Misc (12)

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPRESS OR IMPLIED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON-INFRINGEMENT. BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED TO REPAIR, REPLACEMENT OF OR REFUND FOR THE NON-CONFORMING PRODUCT(S) AT SELLER'S SOLE OPTION. THERE IS NO OBLIGATION TO REPAIR, REPLACE OR REFUND FOR PRODUCTS AS THE RESULT OF (i) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (ii) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (iii) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (iv) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses, or any type of consumption by or application to human or animals.