

CD34 Monoclonal Antibody (QBEND/10)

Product Details

Size	200 µg
Species Reactivity	Human, Non-human primate
Published Species	Rabbit, Human, Mouse
Host/Isotype	Mouse / IgG1
Class	Monoclonal
Type	Antibody
Clone	QBEND/10
Conjugate	Unconjugated
Immunogen	Human endothelial cell membrane vesicles
Form	Liquid
Concentration	1 mg/mL
Purification	Protein A
Storage buffer	PBS
Contains	0.09% sodium azide
Storage conditions	4° C, do not freeze
RRID	AB_2538403

Applications	Tested Dilution	Publications
Western Blot (WB)	Assay-dependent	-
Immunohistochemistry (IHC)	-	59 Publications
Immunohistochemistry (Paraffin) (IHC (P))	1:500-1:1,000	2 Publications
Immunohistochemistry (Frozen) (IHC (F))	Assay-dependent	-
Immunocytochemistry (ICC/IF)	Assay-dependent	3 Publications
Flow Cytometry (Flow)	1:100-1:200	-
Immunoprecipitation (IP)	Assay-dependent	-
Immunomicroscopy (IM)	Assay-dependent	-
Miscellaneous PubMed (Misc)	-	1 Publication


Product Specific Information

This antibody does not react with bovine, dog, rat or sheep.

For FACS analysis, use 10 µL of the suggested working dilution to label 1×10^6 cells in 100 µL. A suggested positive control for immunohistochemical applications is tonsil or bone marrow. This product does not require antigen retrieval using heat treatment or protein digestion. However staining of paraffin-embedded, formalin-fixed tissues may be enhanced by enzyme pre-treatment using pronase or heat treatment using citrate buffer.

Mouse anti Human CD34 antibody, clone QBEND/10 recognizes the human CD34 antigen, also known as Hematopoietic progenitor cell antigen CD34.

Product Images For CD34 Monoclonal Antibody (QBEND/10)



CD34 Antibody (MA5-16926) in IHC (P)
Immunohistochemical analysis of CD34 in paraffin processed placenta. Samples were incubated in CD34 Monoclonal antibody (Product # MA5-16926).



Immunohistochemistry (59)

<p>PloS one</p> <p>CK19 and Glypican 3 Expression Profiling in the Prognostic Indication for Patients with HCC after Surgical Resection.</p> <p>"MA5-16926 was used in immunohistochemistry to find a prognostic indicator for patients with HCC after surgical resection by profiling CK19 and glypican 3 expression"</p> <p>Authors: Feng J,Zhu R,Chang C,Yu L,Cao F,Zhu G,Chen F,Xia H,Lv F,Zhang S,Sun L</p>	<p>Year</p> <p>2016</p>
	<p>Dilution</p> <p>1:100</p>
<p>Romanian journal of morphology and embryology = Revue roumaine de morphologie et embryologie</p> <p>Immunohistochemical study of Ki67, CD34 and p53 expression in human tooth buds.</p> <p>"MA5-16926 was used in immunohistochemistry to study the developmental time course of Ki67, CD34 and p53 expression in human tooth buds"</p> <p>Authors: Muica Nagy-Bota MC,Pap Z,Denes L,Ghizdav A,Brînzaniuc K,Lup Coarc AS,Chibelea Cire-Mrginean M,Pcurar M,Pávai Z</p>	<p>Year</p> <p>2015</p>
	<p>Species</p> <p>Human</p>
	<p>Dilution</p> <p>1:200</p>

[View more IHC references on thermofisher.cn](#)

Immunohistochemistry (Paraffin) (2)

<p>Pathology, research and practice</p> <p>Investigation of the origin of stromal and endothelial cells at the desmoplastic interface in xenograft tumor in mice.</p> <p>"MA5-16926 was used in immunohistochemistry - paraffin section to determine the desmoplastic interface in xenograft tumor in mice comprised of stromal and endothelial cells"</p> <p>Authors: Jung M,Ryu YJ,Kang G</p>	<p>Year</p> <p>2015</p>
	<p>Dilution</p> <p>1:100</p>
<p>International journal of surgical pathology</p> <p>Epstein-Barr virus-associated adrenal smooth muscle tumors and disseminated diffuse large B-cell lymphoma in a child with common variable immunodeficiency: a case report and review of the literature.</p> <p>"MA516926 was used in immunohistochemistry - paraffin section to investigate the clinical and pathological features in a girl with common variable immunodeficiency"</p> <p>Authors: Petrilli G,Lorenzi L,Paracchini R,Ubiali A,Schumacher RF,Cabassa P,Facchetti F</p>	<p>Year</p> <p>2014</p>
	<p>Species</p> <p>Human</p>

More applications with references on thermofisher.cn

- ICC/IF (3)
- Misc (1)

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPRESS OR IMPLIED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT. BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED TO REPAIR, REPLACEMENT OF OR REFUND FOR THE NON-CONFORMING PRODUCT(S) AT SELLER'S SOLE OPTION. THERE IS NO OBLIGATION TO REPAIR, REPLACE OR REFUND FOR PRODUCTS AS THE RESULT OF (i) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (ii) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (iii) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (iv) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses, or any type of consumption by or application to human or animals.