




PREPL Recombinant Rabbit Monoclonal Antibody (JE65-64)

Product Details	
Size	100 μL
Species Reactivity	Human, Mouse, Rat
Host/Isotype	Rabbit / IgG
Expression system	HEK293 cells
Class	Recombinant Monoclonal
Туре	Antibody
Clone	JE65-64
Conjugate	Unconjugated
Immunogen	Recombinant protein within human PREPL aa 621-720/727.
Form	Liquid
Concentration	1 mg/mL
Purification	Protein A
Storage buffer	TBS, pH 7.4, with 40% glycerol, 0.05% BSA
Contains	0.05% sodium azide
Storage conditions	Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.


Applications	Tested Dilution	Publications
Western Blot (WB)	1:500	-
Immunohistochemistry (Paraffin) (IHC (P))	1:400	-

Product Images For PREPL Recombinant Rabbit Monoclonal Antibody (JE65-64)




PREPL Antibody (MA5-44739) in IHC (P)

Immunohistochemistry analysis of PREPL in paraffin-embedded mouse hippocampus tissue. The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH2O and PBS, and then probed with PREPL Monoclonal antibody (Product # MA5-44739) using a dilution of 1:400 for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



PREPL Antibody (MA5-44739) in IHC (P)

Immunohistochemistry analysis of PREPL in paraffin-embedded rat hippocampus tissue. The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH2O and PBS, and then probed with PREPL Monoclonal antibody (Product # MA5-44739) using a dilution of 1:400 for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



PREPL Antibody (MA5-44739) in IHC (P)

Immunohistochemistry analysis of PREPL in paraffin-embedded mouse brain tissue. The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH2O and PBS, and then probed with PREPL Monoclonal antibody (Product # MA5-44739) using a dilution of 1:400 for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

View more figures on thermofisher.cn

For Research Use Only, Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production aspecifications and/or accompanying package inserts ("Documentation"), No claim of studied to prove a paper and the product is usually conformed as a paper and the product is usually conformed as a paper and the product is usually conformed as a paper and the product is usually conformed as a paper and the product is usually conformed as a paper and the product is usually conformed as a paper and the product is usually conformed as a paper and the product is usually conformed as a paper and the product is usually conformed as a paper and the product is usually conformed as a paper and the product is usually conformed as a paper and the product is usually conformed as a paper and the product is usually conformed as a paper and the product is usually conformed as a paper and the product is usually conformed as a paper and the product is usually conformed as a paper and the product is usually conformed as a paper and the product is usually conformed and the product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, or vivo or in vivo therapeutic uses.