


# Butyryl-Histone H2B (Lys5) Polyclonal Antibody

Product Details	
Size	100 µL
Species Reactivity	Human
Host/Isotype	Rabbit / IgG
Class	Polyclonal
Type	Antibody
Conjugate	Unconjugated
Immunogen	Peptide sequence around site of Butyryl-Lys (5) derived from Human Histone H2B type 1-C/E/F/G/I (2-13aa)
Form	Liquid
Concentration	0.18 mg/mL
Purification	Antigen affinity chromatography
Storage buffer	PBS, pH 7.4, with 50% glycerol
Contains	0.03% ProClin 300
Storage conditions	-20°C or -80°C if preferred
RRID	AB_2867207


Applications	Tested Dilution	Publications
Western Blot (WB)	1:100-1:1,000	-
Immunocytochemistry (ICC/IF)	1:10-1:100	-
ELISA (ELISA)	Assay-dependent	-
Immunoprecipitation (IP)	1:200-1:2,000	-
ChIP assay (ChIP)	Assay-dependent	-

Product Images For Butyryl-Histone H2B (Lys5) Polyclonal Antibody




Butyryl-Histone H2B (Lys5) Antibody (PA5-112472) in WB

Western Blot analysis of Butyryl-Histone H2B (Lys5) in multiple sample types using a Butyryl-Histone H2B (Lys5) polyclonal antibody (Product # PA5-112472) at a dilution of 1:100. A goat anti rabbit polyclonal secondary antibody was used at a dilution of 1:50,000. Sample types that were tested include: Hela whole cell lysate, 293 whole cell lysate, A549 whole cell lysate, HepG2 whole cell lysate; Untreated (-) or treated (+) with 30mM sodium butyrate for 4h .



Butyryl-Histone H2B (Lys5) Antibody (PA5-112472) in ICC/IF

Immunocytochemical analysis of Butyryl-Histone H2B (Lys5) in Hela cells using a Polyclonal antibody (Product # PA5-112472) at a dilution of 1:15. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated Goat Anti-Rabbit IgG(H+L). Cells were also counter-stained with DAPI.



Butyryl-Histone H2B (Lys5) Antibody (PA5-112472) in ChIP

Chromatin Immunoprecipitation analysis of Butyryl-Histone H2B (Lys5) in 10<sup>6</sup> Hela cells using a Butyryl-Histone H2B (Lys5) polyclonal antibody (Product #PA5-112472). Hela cells were treated with Micrococcal Nuclease, sonicated, and immunoprecipitated with 5 µg anti-HIST1H2BC or a control normal rabbit IgG. The resulting ChIP DNA was quantified using real-time PCR with primers against the beta-Globin promoter.

View more figures on [thermofisher.cn](http://thermofisher.cn)

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPRESS OR IMPLIED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON-INFRINGEMENT. BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED TO REPAIR, REPLACEMENT OF OR REFUND FOR THE NON-CONFORMING PRODUCT(S) AT SELLER'S SOLE OPTION. THERE IS NO OBLIGATION TO REPAIR, REPLACE OR REFUND FOR PRODUCTS AS THE RESULT OF (i) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (ii) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (iii) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (iv) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses, or any type of consumption by or application to human or animals.