

# Caspase 9 (Cleaved Asp315) Polyclonal Antibody

## Product Details

Size	100 µL
Species Reactivity	Human
Published Species	Human
Host/Isotype	Rabbit / IgG
Class	Polyclonal
Type	Antibody
Conjugate	Unconjugated
Immunogen	Synthetic peptide corresponding to residues surrounding Asp315 of human caspase-9
Form	Liquid
Concentration	6 µg/mL
Purification	Antigen affinity chromatography
Storage buffer	0.01M HEPES, pH 7.5, with 0.15M NaCl, 100µg/mL BSA, 50% glycerol
Contains	no preservative
Storage conditions	-20°C
RRID	AB_10979038

Applications	Tested Dilution	Publications
Western Blot (WB)	1:1,000	2 Publications
Immunocytochemistry (ICC/IF)	1:100	4 Publications
Immunoprecipitation (IP)	1:100	-

## Product Specific Information


It is not recommended to aliquot this antibody.

This antibody was originally validated as part of a Thermo Scientific Cellomics High Content Screening Kit. The antibody sold separately may have slightly different performance and may need to be further optimized for the best results.

Product Images For Caspase 9 (Cleaved Asp315) Polyclonal Antibody


Caspase 9 (Cleaved Asp315) Antibody (PA5-17913) in ICC/IF

Immunofluorescence analysis of Caspase 9 (Cleaved Asp315) was performed using 70% confluent log phase HeLa cells treated with 1µM of Staurosporine for 3 hours. The cells were fixed with 4% paraformaldehyde for 10 minutes, permeabilized with 0.1% Triton™ X-100 for 15 minutes, and blocked with 1% BSA for 1 hour at room temperature. The cells were labeled with Caspase 9 (Cleaved Asp315) Rabbit Polyclonal Antibody (Product # PA5-17913) at 1:100 dilution in 0.1% BSA, incubated at 4 degree Celsius overnight and then labeled with Goat anti-Rabbit IgG (Heavy Chain) Superclonal™ Secondary Antibody, Alexa Fluor® 488 conjugate (Product # A27034) at a dilution of 1:2000 for 45 minutes at room temperature (Panel a: green). Nuclei (Panel b: blue) were stained with SlowFade® Gold Antifade Mountant with DAPI (Product # S36938). F-actin (Panel c: red) was stained with Rhodamine Phalloidin (Product # R415, 1:300). Panel d represents the merged image showing cytoplasmic and nuclear localization. Panel e shows untreated cells with no signal. Panel f represents control cells with no primary antibody to assess background. The images were captured at 60X magnification.




Caspase 9 (Cleaved Asp315) Antibody (PA5-17913)

Altered expression of target protein upon cell treatment demonstrates antibody specificity. Immunofluorescence analysis of Caspase 9 (Cleaved Asp315) using Anti-Caspase 9 (Cleaved Asp315) Rabbit polyclonal Antibody (Product # PA5-17913) shows expression of Caspase 9 (Cleaved Asp315) in HeLa cell line upon Staurosporine treatment. {TM}



Caspase 9 (Cleaved Asp315) Antibody (PA5-17913) in ICC/IF

Immunofluorescent analysis of Cleaved Caspase-9 (green) in HeLa cells either left untreated (left lane) or treated with 1µM Staurosporine for 3 hours. Formalin fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 10 minutes at room temperature and blocked with 1% Blocker BSA (Product # 37525) for 15 minutes at room temperature. Cells were probed with a Cleaved Caspase-9 polyclonal antibody (Product # PA5-17913) at a dilution of 1:100 for at least 1 hour at room temperature, washed with PBS, and incubated with DyLight 488 goat anti-rabbit IgG secondary antibody (Product # 35552) at a dilution of 1:400 for 30 minutes at room temperature. F-Actin (red) was stained with Dylight 554 Phalloidin (Product # 21834) and nuclei (blue) were stained with Hoechst 33342 dye (Product # 62249). Images were taken on a Thermo Scientific ArrayScan or a ToxInsight Instrument at 20X magnification.



View more figures on [thermofisher.cn](http://thermofisher.cn)

## Western Blot (2)

<p><b>Molecular medicine reports</b></p> <p><b>miR302d3p regulates the viability, migration and apoptosis of breast cancer cells through regulating the TMBIM6mediated ERK signaling pathway.</b></p> <p>"Published figure using Caspase 9 (Cleaved Asp315) polyclonal antibody (Product # PA5-17913) in Western Blot"</p> <p>Authors: Liao Y,Qiu Z,Bai L</p>	<p><b>Year</b> 2021</p> <p><b>Species</b> Human</p> <p><b>Dilution</b> 1:1000</p>
<p><b>Oncotarget</b></p> <p><b>Inhibition of the prolyl isomerase Pin1 enhances the ability of sorafenib to induce cell death and inhibit tumor growth in hepatocellular carcinoma.</b></p> <p>"Published figure using Caspase 9 (Cleaved Asp315) polyclonal antibody (Product # PA5-17913) in Western Blot"</p> <p>Authors: Zheng M,Xu H,Liao XH,Chen CP,Zhang AL,Lu W,Wang L,Yang D,Wang J,Liu H,Zhou XZ,Lu KP</p>	<p><b>Year</b> 2017</p>

## Immunocytochemistry (4)

<p><b>Scientific reports</b></p> <p><b>Purine nucleoside phosphorylase deficiency induces p53-mediated intrinsic apoptosis in human induced pluripotent stem cell-derived neurons.</b></p> <p>"PA5-17913 was used in Immunocytochemistry-immunoflourescence to establish a potential mechanism for the neurological defects observed in PNP-deficient patients and reinforce the critical role that PNP has for neuronal viability."</p> <p>Authors: Tsui M,Biro J,Chan J,Min W,Dobbs K,Notarangelo LD,Grunebaum E</p>	<p><b>Year</b> 2022</p> <p><b>Species</b> Human</p>
<p><b>Marine drugs</b></p> <p><b>Stingray Venom Proteins: Mechanisms of Action Revealed Using a Novel Network Pharmacology Approach.</b></p> <p>"PA5-17913 was used in Immunocytochemistry to develop a novel network pharmacology approach based on multi-omics functional data integration to predict how stingray venom disrupts the physiological systems of target animals."</p> <p>Authors: Kirchhoff KN,Billion A,Voolstra CR,Kremb S,Wilke T,Vilcinskas A</p>	<p><b>Year</b> 2021</p> <p><b>Species</b> Human</p>

[View more ICC/IF references on thermofisher.cn](#)

## More applications with references on thermofisher.cn

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPRESS OR IMPLIED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT. BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED TO REPAIR, REPLACEMENT OF OR REFUND FOR THE NON-CONFORMING PRODUCT(S) AT SELLER'S SOLE OPTION. THERE IS NO OBLIGATION TO REPAIR, REPLACE OR REFUND FOR PRODUCTS AS THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses, or any type of consumption by or application to human or animals.