Performance guarenteed'

### **BCL7A Polyclonal Antibody**

#### **Product Details**

Size	100 µg
Species Reactivity	Human
Host/Isotype	Goat / IgG
Class	Polyclonal
Туре	Antibody
Conjugate	Unconjugated
Immunogen	Synthetic peptide sequence (MKLEASQQNSEEM) corresponding to the C-terminus amino acids of BCL7A
Form	Liquid
Concentration	0.5 mg/mL
Purification	Ammonium sulfate precipitation
Storage buffer	TBS, pH 7.3, with 0.5% BSA
Contains	0.02% sodium azide
Storage conditions	-20° C, Avoid Freeze/Thaw Cycles
RRID	AB_10977865

Applications	Tested Dilution	Publications
Western Blot (WB)	1-3 µg/mL	-
Immunocytochemistry (ICC/IF)	10 μg/mL	-
Flow Cytometry (Flow)	10 μg/mL	-


#### **Product Specific Information**


This antibody is predicted to react with bovine, canine, mouse, porcine and rat based on sequence homology.

This antibody is tested in Peptide ELISA: antibody detection limit dilution 4,000.

#### Product Images For BCL7A Polyclonal Antibody

1





contro

#### BCL7A Antibody (PA5-17933) in ICC/IF

Immunocytochemistry analysis of BCL7A using BCL7A Polyclonal Antibody (Product # PA5-17933) in paraformaldehyde fixed U2OS cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10 µg/mL) followed by Alexa Fluor 488 secondary antibody (2 µg/mL), showing nuclear staining. Actin filaments were stained with phalloidin (red) and the nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10 µg/mL) followed by Alexa Fluor 488 secondary antibody (2 µg/mL).



Western blot analysis of BCL7A using BCL7A Polyclonal Antibody (Product # PA5-17933) (1  $\mu$ g/mL) in staining of Human Lymph Node lysate (35  $\mu$ g protein in RIPA buffer). Detected by chemiluminescence.

# 150kDa 100kDa 75kDa


250kDa

50kDa

### 37kDa

25kDa

## 20kDa



#### BCL7A Antibody (PA5-17933) in Flow

Flow cytometric analysis of BCL7A in Daudi cells using a polyclonal antibody (Product #PA5-17933). Daudi cells (blue line) were paraformaldehyde fixed and permeabilized with 0.5% Triton. The primary antibody was incubated for one hour (10  $\mu$ g/mL) followed by an Alexa Fluor 488 secondary antibody (1  $\mu$ g/mL). IgG control: Unimmunized goat IgG (black line) followed by an Alexa Fluor 488 secondary antibody.

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty ger from date of simple time the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample furnished to Buyer is morely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample furnished. To SUBJECT BURENT, EXPRESS 0R INNERINES, EXPRESS 0R INNERINES, EXPRESS 0R INNERINES, EXPRESS 0R INNERINE, EXPRESS 0