



FGF17 Polyclonal Antibody

Product Details		
Size	100 μg	
Species Reactivity	Human	
Host/Isotype	Goat / IgG	
Class	Polyclonal	
Туре	Antibody	
Conjugate	Unconjugated	
Immunogen	E. coli-derived recombinant human FGF-17 Thr23-Thr216	
Form	Lyophilized	
Concentration	0.2 mg/mL	
Purification	Antigen affinity chromatography	
Storage buffer	PBS with 5% trehalose	
Contains	No Preservative	
Storage conditions	-20° C, Avoid Freeze/Thaw Cycles	
RRID	AB_2610199	

Applications	Tested Dilution	Publications
Western Blot (WB)	0.1 μg/mL	-
Neutralization (Neu)	6-24 μg/mL	-


Product Specific Information

In direct ELISAs and Western blots, approximately 5% cross-reactivity with recombinant human (rh) FGF-8b and recombinant mouse FGF-8c is observed and less than 2% cross-reactivity with rhFGF acidic, rhFGF basic, rhFGF-4, rhFGF-5, rhFGF-6, rhFGF-7, rhFGF-10, and rhFGF-18 is observed.

Reconstitute at 0.2 mg/mL in sterile PBS.


Endoxin level is <0.10 EU per 1 μg of the antibody by the LAL method.

Product Images For FGF17 Polyclonal Antibody



FGF17 Antibody (PA5-46934) in Neu

Neutralization of FGF17 in NR6R-3T3 mouse fibroblast cell line. Samples were incubated in FGF17 polyclonal antibody (Product # PA5-46934). Recombinant Human FGF-17 stimulates proliferation in the NR6R-3T3 mouse fibroblast cell line in a dose-dependent manner (orange line). Proliferation elicited by Recombinant Human FGF-17 (100 ng/mL) is neutralized (green line) by increasing concentrations of Goat Anti-Human FGF-17 Antigen Affinity-purified Polyclonal Antibody. The ND50 is typically 6-24 $\mu g/mL$ in the presence of heparin (1 $\mu g/mL$).



FGF17 Antibody (PA5-46934)

Neutralization antibody testing demonstrates the specificity of an antibody through a correlation between antibody binding and the activity of the target. Neutralization of human FGF-17 is shown by decrease in CPM (measure of proliferation) with increasing concentrations of FGF-17 polyclonal antibody (PA5-46934). {Neu}

For Research Use Only, Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product is pecifications in effect at the time of sale, as set forth in the Production occumentation.

In the Product is pecification and/or accompanying package inserts ("Documentation"), No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPRESS OR IMPLED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICIDATE OR NON INFRINGEMENT.

BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURINGS THE WARRANTY PERFOLO IS LIMITED TO REPAIR, REPLACE OR REFUND FOR THE NON-CONFORMING PRODUCTS) AT SELECTION. THERE IS NO. LO POTION. TH