



AP2M1 Polyclonal Antibody


| Product Details | | |
|------------------------|---|--|
| Size | 100 μg | |
| Species Reactivity | Human, Mouse, Rat | |
| Host/Isotype | Rabbit / IgG | |
| Class | Polyclonal | |
| Туре | Antibody | |
| Conjugate | Unconjugated | |
| Immunogen | A synthetic peptide corresponding to a sequence at the C-terminus of human AP2M1 (399-435aa LKVRYLKVFEPKLNYSDHDVIKWVRYIGRSGIYETRC). | |
| Form | Lyophilized | |
| Concentration | 500 μg/mL | |
| Purification | Affinity chromatography | |
| Storage buffer | PBS with 4mg trehalose | |
| Contains | 0.05mg sodium azide | |
| Storage conditions | Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles. | |
| RRID | AB_2807193 | |

| Applications | Tested Dilution | Publications |
|-------------------|-----------------|--------------|
| Western Blot (WB) | 0.1-0.5 μg/mL | - |

Product Specific Information

Reconstitute with 0.2 mL of distilled water to yield a concentration of 500 µg/mL.

Product Images For AP2M1 Polyclonal Antibody



AP2M1 Antibody (PA5-95390) in WB

Western blot analysis of AP2M1 in, Lane 1: human U2OS whole cell lysates, Lane 2: human MCF-7 whole cell lysates, Lane 3: rat kidney tissue lysates, . Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50 µg of sample under reducing conditions. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. The membrane was blocked with 5% Non-fat Milk/ TBS for 1. 5 hour at RT. The membrane was incubated with AP2M1 Polyclonal Antibody (Product # PA5-95390) at 0.5 g/mL overnight at 4°C, then washed with TBS-0. 1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1. 5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit. A specific band was detected for AP2M1 at approximately 50KD. The expected band size for AP2M1 is at 50KD.

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization, Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation specifications and/or accompanying package instentiation?). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid not hye house duby properly trained individuals. Unless otherwise stated in the Documentation, this warranty provided herein is valid not herein to yet properly trained individuals. Unless otherwise stated in the Universal individuals. Unless otherwise stated in dividuals. Unless otherwise warranty provided herein is valid not herein to yet properly trained individuals. Unless otherwise stated individuals. Unless otherwise stated individuals. Unless otherwise warranty provided herein is valid not be used for any other product is made or sample. No OTHER WARRANTIES, EXPRESS OR IMPLED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT.

BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS OR NON INFRINGEMENT.

BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS OR NON INFRINGEMENT.

BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS SOLD AND THERE IS NO DELIGATION TO REPAIR, REPLACE OR REFUND FOR THE NON-CONFORMING PRODUCTS; AS THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, or vivo or in vivo therapeutic uses,