



NOVA1 Polyclonal Antibody


Product Details		
Size	100 μg	
Species Reactivity	Human, Mouse, Rat	
Host/Isotype	Rabbit / IgG	
Class	Polyclonal	
Туре	Antibody	
Conjugate	Unconjugated	
Immunogen	E.coli-derived human Nova1 recombinant protein (Position: T416-V509).	
Form	Lyophilized	
Concentration	500 μg/mL	
Purification	Affinity chromatography	
Storage buffer	PBS with 4mg trehalose	
Contains	0.05mg sodium azide	
Storage conditions	Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.	
RRID	AB_2807373	

Applications	Tested Dilution	Publications
Western Blot (WB)	0.1-0.5 μg/mL	-
Immunohistochemistry (Paraffin) (IHC (P))	0.5-1 μg/mL	-
Immunohistochemistry (Frozen) (IHC (F))	0.5-1 μg/mL	-
Immunocytochemistry (ICC/IF)	2 μg/mL	-
ELISA (ELISA)	0.1-0.5 μg/mL	-

Product Specific Information


Reconstitute with 0.2 mL of distilled water to yield a concentration of 500 $\mu g/mL$.

Product Images For NOVA1 Polyclonal Antibody




NOVA1 Antibody (PA5-95571) in WB

Western blot analysis of NOVA1 in Lane 1: human U-87MG whole cell lysate, Lane 2: human Caco-2 whole cell lysate. Electrophoresis was performed with 5-20% SDS-PAGE gel (70V, Stacking gel; 90V Resolving gel, Time: 2-3 hours), transferred to a nitrocellulose membrane and blocked using 5% Non-fat Milk/TBS (1.5 hrs at room temperature). Samples were incubated with NOVA1 polyclonal antibody (Product # PA5-95571) using a 0.5 μ g/mL dilution, followed by a goat anti-rabbit IgG-HRP at a dilution of 1:10,000, and developed with enhanced chemiluminescence (ECL).



NOVA1 Antibody (PA5-95571) in ICC/IF

Immunofluorescence analysis of NOVA1 in A549 cell. Antigen retrieval was performed on the tissue using IHC enzyme antigen retrieval reagent and blocked with 10% goat serum. Samples were incubated with NOVA1 polyclonal antibody (Product # PA5-95571) at a 2 μ g/mL dilution, followed by 488 conjugated goat anti-rabbit IgG using a 1:100 dilution.



NOVA1 Antibody (PA5-95571) in IHC (P)

Immunohistochemistry analysis of NOVA1 in paraffin-embedded human glioma tissue. Antigen retrieval was performed on the tissue using citrate buffer (pH 6, 20 min) and blocked with 10% goat serum. Samples were incubated with NOVA1 polyclonal antibody (Product # PA5-95571) at a 1 μ g/mL dilution, followed by biotinylated goat anti-rabbit IgG (30 min, 37°C), and developed with Strepavidin-Biotin-Complex and DAB.

View more figures on thermofisher.cn

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPERESS OR IMPLEA REPRESS OR IMPLEA REGRANTED INITY, IMPLIED WARRANTIES, EXPERESS OR IMPLEA REPRESS OR IMPLEAD IN INFINITY ("INFINITY PROPOSE"). AND ANY PARTICULAR VARRANTIES, EXPERESS OR IMPLEAD IN INFINITY PROPOSED, OR NON INFINITY PROPOSED IN INFINITY PROPOSED I