Performance guarenteed

EVI1 Polyclonal Antibody

Product Details

Size	100 µL
Species Reactivity	Human, Rat
Host/Isotype	Rabbit / IgG
Class	Polyclonal
Туре	Antibody
Conjugate	Unconjugated
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 1-190 of human MECOM (NP_0049822)
Form	Liquid
Concentration	0.34 mg/mL
Purification	Affinity Chromatography
Storage buffer	PBS, pH 7.3, with 50% glycerol
Contains	0.02% sodium azide
Storage conditions	-20° C, Avoid Freeze/Thaw Cycles
RRID	AB_2808186

Applications	Tested Dilution	Publications
Western Blot (WB)	1:500-1:2,000	-


Product Specific Information

Positive Samples: Mouse kidney, Mouse heart; Cellular Location: Nucleus, Nucleus speckle

Immunogen sequence: MRSKGRARKL ATNNECVYGN YPEIPLEEMP DADGVASTPS LNIQEPCSPA TSSEAFTPKE GSPYKAPIYI PDDIPIPAEF ELRESNMPGA GLGIWTKRKI EVGEKFGPYV GEQRSNLKDP SYGWEILDEF YNVKFCIDAS QPDVGSWLKY IRFAGCYDQH NLVACQINDQ IFYRVVADIA PGEELLLFMK


Product Images For EVI1 Polyclonal Antibody

1



EVI1 Antibody (PA5-96384) in WB

Western blot analysis of EVI1 in various lysates. Samples were incubated with EVI1 Polyclonal antibody (Product # PA5-96384) using a dilution of 1:2,500, followed by HRP Goat Anti-Rabbit IgG (H+L) at a dilution of 1:10,000. Lysates /proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 30s.



EVI1 Antibody (PA5-96384) in WB

Western blot analysis of EVI1 in rat kidney. Samples were incubated with EVI1 Polyclonal antibody (Product # PA5-96384) using a dilution of 1:2,500, followed by HRP Goat Anti-Rabbit IgG (H+L) at a dilution of 1:10,000. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 180s.

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by property trained individuals. Unless otherwise stated in the Documentation, this warranty is provided herein is valid only when used by property trained individuals. Unless otherwise stated in the Documentation, this warranty is provided herein is valid only when used by property trained individuals. Unless otherwise stated is anyone other than the Buyer. Any model or sample furnisated to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. No OTHER WARRANTES, EXPRESS OR INNERNICE, EXPRESS OR INNERNICE CREMENT CONCLUSIVE REMEMPT FOR NON-CONFORMING PRODUCTS DURING THE WARRANTED, EXCLUSIVE CREMENT FOR NON-CONFORMING PRODUCTS DURING THE WARRANT PERIOD IS LIMITED TO REPAIR, REPLACEMENT OF OR REFLUND FOR THE NON-CONFORMING PRODUCTS SOLE OFTION. THERE IS NO BUIGATION TO REPAIR, REPLACE OR REFLUND FOR THE NON-CONFORMING PRODUCTS SOLE OFTION. THERE IS NO BUIGATION TO REPAIR, REPLACE OR REFLUND FOR THE NON-CONFORMING PRODUCTS AS THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDILLON OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, are vity or or invito