

## Human CCL20 (MIP-3 alpha) Recombinant Protein

| <b>Product Details</b>  |   |  |  |
|-------------------------|---|--|--|
| Size                    | 20 μg   |  |  |
| Species                 | Human   |  |  |
| Expression system       | E. coli   |  |  |
| Amino acid sequence     | Human CCL20 recombinant protein contains 70 amino acids                             |  |  |
| Molecular weight        | 8 kDa   |  |  |
| Class                   | Recombinant   |  |  |
| Туре                    | Protein   |  |  |
| Purity                  | >98% by SDS-PAGE  |  |  |
| Endotoxin concentration | <0.1 ng/µg  |  |  |
| Activity                | ED50 = 10 - 50 ng/mL; determined by the dose-dependent chemotaxis of human T-cells. |  |  |
| Conjugate               | Unconjugated  |  |  |
| Form                    | Lyophilized   |  |  |
| Purification            | sequential chromatography   |  |  |
| Contains                | no preservative   |  |  |
| Storage conditions      | -20°C   |  |  |

| Applications               | Tested Dilution | Publications |
|----------------------------|-----------------|--------------|
| Western Blot (WB)          | Assay-dependent | -            |
| Immunohistochemistry (IHC) | Assay-dependent | -            |
| ELISA (ELISA)              | Assay-dependent | -            |
| Functional Assay (FN)      | Assay-dependent | -            |
| Control (Ctrl)             | Assay-dependent | -            |

## **Product Specific Information**

Carrier-Free

Reconstitution: We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute recombinant human MIP-3 alpha in sterile, distilled water to a concentration of 0.1-1.0 mg/mL. Further dilutions should be made in tissue culture grade buffered solutions with heat inactivated fetal calf serum or low endotoxin BSA.

Storage: This lyophilized preparation is stable at 2°C to 8°C, but should be kept at -20°C for long term storage, preferably desiccated. Apportion the reconstituted preparation into working aliquots and store at -20°C. Avoid repeated freeze/thaw cycles.

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production obcumentation, specifications and/or accompanying package inserts ("Documentation,"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is imited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPRESS OR IMPLED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NOI INFRINGEMENT.

BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED TO REPAIR, REPLACE OR REFUND FOR THE NON-CONFORMING PRODUCTS, AT SELLER'S SOLE OPTION. THERE IS NO OBLIGATION TO REPAIR, REPLACE OR REFUND FOR PRODUCTS AS THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISSUE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, or vivy for efforts.